	Research Ethics	Target no	Recruitment	Did the trial Recruit to
NAME OF TRIAL	No (REC)	Patients	Deadline Trial S	tatus Target?
A long term monitoring study to evaluate the persistence of direct acting antiviral (DAA) treatment resistant mutations or the durability of sustained virological response (SVR) in patients treated with DAA containing regimens for chronic hepatitis C infection (CHC) A MULTICENTER, PHASE III, OPEN-LABEL, RANDOMIZED STUDY IN PREVIOUSLY	11/LO/0278	6	30/09/2012 Open	Still Open
UNTREATED PATIENTS WITH ADVANCED INDOLENT NON-HODGKIN'S LYMPHOMA EVALUATING THE BENEFIT OF GA101 (RO5072759) PLUS CHEMOTHERAPY COMPARED WITH RITUXIMAB PLUS CHEMOTHERAPY FOLLOWED BY GA101	11/NW/0298	6	30/06/2013 Open	Still Open
A Multi-Centre 3-Year Follow-Up study to Assess the Durability of Sustained Virologic Resonse in Alisporivir-Treated Chronic Hepatitis C Patients	12/SC/0185	1	23/03/2015 Open	Still Open
A new vaccine against meningitis groupB: A second follow on study	12/SC/0384	10		Still Open
A Phase 3 Evaluation of Daclatasvir in Combination with Peg-interferon Lambda-1a and Ribavirin (RBV) or Telaprevir in Combination with Peginterferon Alfa-2a and RBV in Patients with Chronic Hepatitis C Genotype				
1b who are Treatment Naïve or Prior Relapsers to Alfa/RBV Therapy	13/LO/0206	4	26/09/2013 Open	Still Open
A Phase 3 Study with Asunaprevir and Daclatasvir (DUAL) for Null or Partial Responders to Peginterferon Alfa and Ribavirin (P/R), Intolerant or Ineligible to P/R Subjects and Treatment-Naive Subjects with Chronic Hepatitis C				
Genotype 1b Infection A Phase 3, Randomised, Double-blind, Placebo-controlled, Parallel-group,	12/LO/0827	5	20/03/2014 Open	Still Open
Multi-centre Study to Evaluate the Safety and Efficacy of Ustekinumab Induction Therapy in Participants with Moderately to Severely Active Crohn?s	11/50/0227	-	15/00/2012 0	Chill On an
Disease A Phase 3, Randomised, Double-blind, Placebo-controlled, Parallel-group,	11/SC/0327	5	15/09/2013 Open	Still Open
Multi-centre Study to Evaluate the Safety and Efficacy of Ustekinumab Maintenance Therapy in Subjects with Moderately to Severely Active Crohn's Disease	11/SC/0329	5	15/09/2013 Open	Still Open
A phase III open-label randomised study, to evaluate the immunogenicity and safety of the concomitant administration of V419 (PRSI) given at 2, 3 and 4 months of age with two types of meningococcal serogroup C conjugate (MCC) vaccines given at 3 and 4 months of age, and followed by the administration at 12 months of age of a combined Haemophilus influenzae type b-MCC				
vaccine.	11/SW/0328	30	31/08/2012 Open	Still Open
A phase III randomised, partially double-blind and placebo-controlled study of BI 207127 in combination with faldaprevir and ribavirin for chronic genotype 1 hepatitis C infection in an extended population of treatment naïve patients that includes those ineligible to receive peginterferon.		5	07/05/2013 Open	Still Open
A PHASE III, DOUBLE-BLIND, PLACEBO-CONTROLLEDSTUDY OF VEMURAFENIB VERSUS VEMURAFENIB PLUS GDC-0973 IN PREVIOUSLY UNTREATED			No date	·
BRAFV600-MUTATION POSITIVE PATIENTS WITH UNRESECTABLE LOCALLY	12/10/1976	c	agreed with Sponsor Open	C+ill Onon
ADVANCED OR METASTATIC MELANOMA A phase III, open-label study of once daily BI 201335 240 mg for 24 weeks in combination with pegylated interferon-? (PegIFN) and ribavirin (RBV) in patients with genotype 1 chronic hepatitis C infection who failed a prior	12/LO/1876		Sponsor Open	Still Open
PegIFN / RBV treatment A Phase III, Randomized, Comparative, Open-label Study of Intravenous Iron	11/SC/0177	1	31/08/2013 Open	Still Open
Isomaltoside 1000 (Monofer?) Administered as Maintenance Therapy by Single or Repeated Bolus Injections in Comparison with Intravenous Iron Sucrose in Subjects with Stage 5 Chronic Kidney Disease on Dialysis Therapy				
(CKD-5D). A Phase II Study of Dasatinib Therapy in Children and Adolescents with Ph+	11/NE/0124	10	31/07/2013 Open	Still Open
Leukemia with Resistance of Inolerance to Imatinib Phase I/II study of Scr/ABi tyrosine kinase inhibitor dasatinib (BMS-354825) in children and adolescents				
with relapsed or refractory leukemia	09/H0405/51	4		Still Open
A Proof of Concept Study for GSK249320 versus placebo in Stroke Patients A Randomised, Double Blind, Parallel Group, Multicentre Phase IIb study to	11/LO/1447	8	31/10/2014 Open	Still Open
complare ticagrelor with clopidogrel treatment on the risk of cardiovascular deat, myocardial infarction and ischeamic stroke in patients with established	12/504/0005	22	14/05/2011	Critt C
Peripheral Artery Disease A randomized, controlled, open-label, parallel-group, multi-center study to compare the effect of Intrathecal Baclofen Therapy (ITB Therapy?) versus Best	12/EM/0395	20	14/06/2014 Open	Still Open
Medical Treatment (BMT) on severe spasticity in post-stroke patients after 6 months active treatment	10/H0803/96	4	31/03/2011 Open	Still Open

A Randomized, Double-Blind, Phase II, Efficacy and Safety Study of MDV3100			
(ASP9785) vs. Bicalutamide in Castrate Men with Metastatic Prostate Cancer	11/LO/0608	5 12/07/2012 Open	Still Open
A study exploring the efficacy and safety of once-daily oral Rivaroxaban (BAY			
59-7939) compared with that of dose-adjusted oral vitamin K antagonist (VKA)			
for the prevention of cardiovascular events in subjects with nonvalvular atrial fibrillation scheduled for cardioversion	12/10/1205	43 30/00/3043 0	Chill On an
This inaction scriedated for cardioversion	12/LO/1285	12 28/09/2013 Open	Still Open
Adjuvant immunotherapy with anti-CTLA-4 monoclonal antibody (ipilimumab)			
versus placebo after complete resection of high-risk Stage III melanoma: A			
randomized, double-blind Phase 3 trial of the EORTC Melanoma Group	08/H0311/120	30 01/06/2013 Open	Still Open
BIA-2093-311: Efficacy and safety of eslicarbazepine acetate (BIA 2-093) as			
monotherapy for patients with newly diagnosed partial-onset seizures: A			
double-blind, randomized, active-controlled, parallel-group, multicenter clinical study	10/H0903/29	6 01/12/2013 Open	Still Open
Clinical Protocol Al444026: An Open-Label Re-treatment Study with Peg-	10/110303/23	0 01/12/2013 Open	Juli Open
Interferon Alfa-2a, Ribavirin and MBS-790052 With or Without BMS-650032			
for subjects with Chronic Hepatitis C	11/SC/0358	4 30/09/2015 Open	Still Open
		No date	
OVATION POST MARKET STUDY Outlier Alideratical Start Conft Surtain	44/10/0022	agreed with	Chill Owner
OVATION POST-MARKET STUDY Ovation Abdominal Stent Graft System	11/LO/0922	15 Sponsor Open	Still Open
Randomised, phase IV, placebo-controlled, comparative study to evaluate the			
efficacy and safety of tapering methotrexate (MTX) dosage versus maintaining			
the dosage in patients with severe active Rheumatoid Arthritis (RA) who have			
demonstrated an inadequate response to prior disease-modifying anti-			
rheumatic drugs (DMARDS) treatment and have initiated RoActemra®	12/55/0176	5 04/40/2040 0	C::!! C
(Tocilizumab, TCZ) in combination with MTX.	12/EE/0176	5 31/12/2013 Open	Still Open
Surgical replacement and transcatheter aortic valve implantation	12/LO/0491	25 31/10/2018 Open	Still Open
Zenith TX2 Low-Profile TAA Endovascular Graft Clinical Study #08-017-03	10/H0808/27	10 30/09/2010 Open	Still Open
A Phase 3 Evaluation of BMS-790052 in Combination with Peg-Tnerferon Alfa-			
2a and Ribavirin in Treatment-Nainve Subjects with Chronic Hepatitis C	12/56/0024	2 20/04/2014 02 27	21/2
Genotypes 1 and 4 A Phase 3, Multi-centre, Randomised, Double-blind, Placebo-controlled,	12/SC/0031	2 30/04/2014 Open	N/A
Parallel-group study of the Efficacy and Safety of Lenalidomide (Revlimid?) as			
Maintenance Therapy for Patients with B-Cell Chronic Lymphocytic Leukaemia			
following Second-line Therapy.	09/H0301/5	5 30/11/2013 Open	N/A
A Phase II single blind, randomized, placebo controlled trial to study the			
efficacy and safety of anti-von Willebrand factor Nanobody administered as			
adjunctive treatment to patients with acquired thrombotic thrombocytopenic	10/40700/60	3 21/05/2012 Open	NI/A
purpura. A Phase II, Multicenter, Open-Label Study Of YM155 Plus Rituximab In	10/H0709/60	3 31/05/2012 Open	N/A
Previously Treated Subjects With CD20-Positive B Cell Non-Hodgkin?s			
Lymphoma Who Are Ineligible For Or Have Previously Received An Autologous			
Stem Cell Transplant	10/H0406/16	5 30/06/2013 Open	N/A
A phase III, randomised, double-blind and placebo controlled study of once		No data	
daily BI 201335, 240 mg for 12 or 24 weeks in combination with pegylated interferon-? and ribavirin in patients with genotype 1 chronic hepatitis C		No date agreed with	
infection who failed a prior PegIFN/RBV treatment	11/SC/0176	5 Sponsor Open	N/A
A Randomized, Double-Blind, Multicenter Study of Denosumab Compared	,, -	- sp.	•
With Zoledronic Acid (Zometa?) in the Treatment of Bone Disease in Subjects			
with Newly Diagnosed Multiple Myeloma	12/LO/1076	5 31/05/2015 Open	N/A
An open Phase I Study of immunization with the recNY-ESO-1 + AS15 Antigen-			
Specific Cancer Immunotherapeutic in patients with NY-ESO-1-positive			
unresectable and progressive metastatic cutaneous melanoma.	10/H0806/106	4 30/12/2011 Open No date	N/A
A multi-centre double-blind parallel-group placebo-controlled study of the efficacy and safety of teriflunomide in patients with relapsing multiple		agreed with	
sclerosis who are treated with interferon-beta	11/H0707/6	6 Sponsor Withdrawn	N/A
		No date	•
Comparison of New Ostomy Barriers and Drainable Pouches to currently		agreed with	
Marketed Ostomy Pouching Systems	12/EE/0078	10 Sponsor Withdrawn	N/A
European Study of POBA versus Cotavance Paclitaxel Coated Balloon for the	11/10/0356	45 20/02/2042 1991	N1 / A
Treatment of Infrapopliteal Lesions in Critical Limb Ischemia	11/LO/0256	15 28/02/2013 Withdrawn	N/A
Randomized, Double-Blind, Controlled Study to Evaluate the Efficacy and			
Safety of the Combination of ABT-450/Ritonavir/ABT-267 (ABT-450/r/ABT-			
267) and ABT-333 With and Without Ribavirin (RBV) in Treatment-Naïve		Closed - In	
Adults with Genotype 1a Chronic Hepatitis C Virus (HCV) Infection (PEARL-IV)	13/LO/0425	5 31/05/2013 Follow Up	Yes
study to assess the incidence of mutations in the tyrosine kinase domain of			
the epidermal growth factor receptor in UK patients with newly diagnosed			
locally advanced or metastatic non-small cell lung cancer and to investigate		No date	
the quality of life of these patients undergoing first-line monotherapy with	10/H0713/75	agreed with Closed - In 10 Sponsor Follow Up	Yes
erlotinib (Tarceva®).			

A multi-center, randomized, parallel-group, rater-blinded study comparing the effectiveness and safety of teriflunomide and interferon beta-1a in			
		Classed In	
I makingka uulkla walagaina gaulkin la salagasia	00/110720/125	Closed - In	Vaa
patients with relapsing multiple sclerosis.	09/H0720/135	4 14/10/2011 Follow Up	Yes
A multi-center, randomized, parallel-group, rater-blinded study comparing		Clared In	
the effectiveness and safety of teriflunomide and interferon beta-1a in	40/114007/0	Closed - In	.,
patients with relapsing multiple sclerosis.	10/H1307/8	4 14/10/2011 Follow Up	Yes
A Phase 3, randomised, double-blind study of the safety and efficacy of			
GSK1349572 plus abacavir/lamivudine fixed-dose combination therapy			
administered once daily compared to Atripla over 96 weeks in HIV-1 infected		Closed - In	
antiretroviral therapy naive adult subjects.	10/H0713/76	2 30/04/2011 Follow Up	Yes
A Phase III Open, Multicentre Study to Investigate the Pharmacokinetics,			
Safety and Efficacy of BPL?s High Purity Factor X in the Treatment of Severe		Closed - In	
and Moderate Factor X Deficiency	09/H1306/82	2 31/12/2012 Follow Up	Yes
A phase III, open-label, multicentre study to evaluate the immunogenicity,			
safety and reactogenicity of revaccination dose of the GlaxoSmithKline			
Biologicals' quadrivalent seasonal influenza candidate vaccine GSK2321138A,			
administered to children who previously participated in study 115345 (FLU D-		Closed - In	
QIV-004 PRI)	12/SW/0237	9 05/07/2013 Follow Up	Yes
Q(V 0041111)	12/34/023/	3 03/07/2013 Follow 0p	163
A DANDOMICED DOUBLE BLIND DOUBLE DLIMMY BLACERO CONTROLLED			
A RANDOMISED, DOUBLE-BLIND, DOUBLE-DUMMY, PLACEBO-CONTROLLED,		No dete	
DOSE-RANGING PHASE II STUDY ASSESSING RANOLAZINE IN THE		No date	
MAINTENANCE OF SINUS RHYTHM AFTER ELECTRICAL CARDIOVERSION IN		agreed with Closed - In	
PATIENTS WITH NON-PERMANENT ATRIAL FIBRILLATION	12/LO/0009	6 Sponsor Follow Up	Yes
An Open-Label, Multicenter Extension Study to Evaluate the Safety and			
Tolerability of Natalizumab Following Re-Initiation of Dosing in Multiple		No date	
Sclerosis Who Have Completed Studies C-1801, C-1802 and a Dosing		agreed with Closed - In	
Suspension Safety Evaluation (STRATA)	06/Q0505/31	Sponsor Follow Up	Yes
· , , , , , , , , , , , , , , , , , , ,	, - =	Closed - In	-
Efficacy Study of a Quadrivalent Influenza Vaccine in Children (FLU D-QIV)	11/SC/0313	25 14/12/2011 Follow Up	Yes
Prospective, Multicenter, Single Arm Feasibility and Safety Study of the	11,00,0010	No date	103
Endologix Fenestrated Stent Graft System for the Endovascular Repair of	/== /0	agreed with Closed - In	
Juxtarenal/Pararenal (JAA/PAA) Aneurysms	11/EE/0219	5 Sponsor Follow Up	Yes
Protocol RTA115450: A two part double blind, randomised, placebo-			
controlled and open-label study to investigate the efficacy, safety and			
tolerability of eltrombopag, a thrombopoietin receptor agonist, in pediatric		No date	
patients with previously treated chronic immume (idopathic)		agreed with Closed - In	
thrombocytopenic purpura (ITP)	12/NW/0134	2 Sponsor Follow Up	Yes
		No date	
To prospectively collect global 'real world' safety and clinical performance		agreed with Closed - In	
data on Endurant Stent Graft System. This is a registry only.	09/H0308/124	10 Sponsor Follow Up	Yes
A double-blind, randomized, placebo-controlled Phase III study to assess the	03/110300/121	10 Sponson 1 show Sp	
efficacy of MAGE-A3 + AS15 ASCI as adjuvant therapy in patients with MAGE-		Closed - In	
	10/110002/20		Na
A3 positive resected stage III melanoma.	10/H0903/39	3 31/08/2016 Follow Up	No
A PHASE 3 STUDY TO EVALUATE THE EFFICACY AND SAFETY OF DOCETAXEL		No date	
AND PREDNISONE WITH OR WITHOUT LENALIDOMIDE IN SUBJECTS WITH		1 11 41 41 1 1	
		agreed with Closed - In	
CASTRATE-RESISTANT PROSTATE CANCER (MAINSAIL)	09/H0720/132	agreed with Closed - In 15 Sponsor Follow Up	No
	09/H0720/132	•	No
CASTRATE-RESISTANT PROSTATE CANCER (MAINSAIL) A Phase 3, Multicentre, Randomised, Open-Label, Parallel-Group Study of the	09/H0720/132	•	No
CASTRATE-RESISTANT PROSTATE CANCER (MAINSAIL) A Phase 3, Multicentre, Randomised, Open-Label, Parallel-Group Study of the Efficacy and Safety of Lenalidomide (Revlimid) Versus Chlorambucil as First-	09/H0720/132	15 Sponsor Follow Up	No
CASTRATE-RESISTANT PROSTATE CANCER (MAINSAIL) A Phase 3, Multicentre, Randomised, Open-Label, Parallel-Group Study of the Efficacy and Safety of Lenalidomide (Revlimid) Versus Chlorambucil as First-Line Therapy for Previously Untreated Elderly Patients with B-Cell Chronic		15 Sponsor Follow Up Closed - In	
CASTRATE-RESISTANT PROSTATE CANCER (MAINSAIL) A Phase 3, Multicentre, Randomised, Open-Label, Parallel-Group Study of the Efficacy and Safety of Lenalidomide (Revlimid) Versus Chlorambucil as First-	09/H0720/132 09/H1107/103	15 Sponsor Follow Up	No No
CASTRATE-RESISTANT PROSTATE CANCER (MAINSAIL) A Phase 3, Multicentre, Randomised, Open-Label, Parallel-Group Study of the Efficacy and Safety of Lenalidomide (Revlimid) Versus Chlorambucil as First-Line Therapy for Previously Untreated Elderly Patients with B-Cell Chronic Lymphocytic Leukaemia. (The Origin Trial)		15 Sponsor Follow Up Closed - In	
CASTRATE-RESISTANT PROSTATE CANCER (MAINSAIL) A Phase 3, Multicentre, Randomised, Open-Label, Parallel-Group Study of the Efficacy and Safety of Lenalidomide (Revlimid) Versus Chlorambucil as First-Line Therapy for Previously Untreated Elderly Patients with B-Cell Chronic Lymphocytic Leukaemia. (The Origin Trial) A Phase 3, Randomised, Double-blind, Placebo-controlled, Parallel-group,		15 Sponsor Follow Up Closed - In	
CASTRATE-RESISTANT PROSTATE CANCER (MAINSAIL) A Phase 3, Multicentre, Randomised, Open-Label, Parallel-Group Study of the Efficacy and Safety of Lenalidomide (Revlimid) Versus Chlorambucil as First-Line Therapy for Previously Untreated Elderly Patients with B-Cell Chronic Lymphocytic Leukaemia. (The Origin Trial) A Phase 3, Randomised, Double-blind, Placebo-controlled, Parallel-group, Multi-centre Study to Evaluate the Safety and Efficacy of Ustekinumab		15 Sponsor Follow Up Closed - In 5 31/05/2011 Follow Up	
CASTRATE-RESISTANT PROSTATE CANCER (MAINSAIL) A Phase 3, Multicentre, Randomised, Open-Label, Parallel-Group Study of the Efficacy and Safety of Lenalidomide (Revlimid) Versus Chlorambucil as First-Line Therapy for Previously Untreated Elderly Patients with B-Cell Chronic Lymphocytic Leukaemia. (The Origin Trial) A Phase 3, Randomised, Double-blind, Placebo-controlled, Parallel-group, Multi-centre Study to Evaluate the Safety and Efficacy of Ustekinumab Induction Therapy in Particiapnts with Moderately to Severely Active Crohn?s	09/H1107/103	15 Sponsor Follow Up Closed - In 5 31/05/2011 Follow Up Closed - In	No
CASTRATE-RESISTANT PROSTATE CANCER (MAINSAIL) A Phase 3, Multicentre, Randomised, Open-Label, Parallel-Group Study of the Efficacy and Safety of Lenalidomide (Revlimid) Versus Chlorambucil as First-Line Therapy for Previously Untreated Elderly Patients with B-Cell Chronic Lymphocytic Leukaemia. (The Origin Trial) A Phase 3, Randomised, Double-blind, Placebo-controlled, Parallel-group, Multi-centre Study to Evaluate the Safety and Efficacy of Ustekinumab		15 Sponsor Follow Up Closed - In 5 31/05/2011 Follow Up	
CASTRATE-RESISTANT PROSTATE CANCER (MAINSAIL) A Phase 3, Multicentre, Randomised, Open-Label, Parallel-Group Study of the Efficacy and Safety of Lenalidomide (Revlimid) Versus Chlorambucil as First-Line Therapy for Previously Untreated Elderly Patients with B-Cell Chronic Lymphocytic Leukaemia. (The Origin Trial) A Phase 3, Randomised, Double-blind, Placebo-controlled, Parallel-group, Multi-centre Study to Evaluate the Safety and Efficacy of Ustekinumab Induction Therapy in Particiapnts with Moderately to Severely Active Crohn?s	09/H1107/103	15 Sponsor Follow Up Closed - In 5 31/05/2011 Follow Up Closed - In	No
CASTRATE-RESISTANT PROSTATE CANCER (MAINSAIL) A Phase 3, Multicentre, Randomised, Open-Label, Parallel-Group Study of the Efficacy and Safety of Lenalidomide (Revlimid) Versus Chlorambucil as First-Line Therapy for Previously Untreated Elderly Patients with B-Cell Chronic Lymphocytic Leukaemia. (The Origin Trial) A Phase 3, Randomised, Double-blind, Placebo-controlled, Parallel-group, Multi-centre Study to Evaluate the Safety and Efficacy of Ustekinumab Induction Therapy in Particiapnts with Moderately to Severely Active Crohn?s	09/H1107/103	15 Sponsor Follow Up Closed - In 5 31/05/2011 Follow Up Closed - In	No
CASTRATE-RESISTANT PROSTATE CANCER (MAINSAIL) A Phase 3, Multicentre, Randomised, Open-Label, Parallel-Group Study of the Efficacy and Safety of Lenalidomide (Revlimid) Versus Chlorambucil as First-Line Therapy for Previously Untreated Elderly Patients with B-Cell Chronic Lymphocytic Leukaemia. (The Origin Trial) A Phase 3, Randomised, Double-blind, Placebo-controlled, Parallel-group, Multi-centre Study to Evaluate the Safety and Efficacy of Ustekinumab Induction Therapy in Particiapnts with Moderately to Severely Active Crohn?s Disease Who Have Failed or Are Intolerant to TNF Antagonist Therapy	09/H1107/103	15 Sponsor Follow Up Closed - In 5 31/05/2011 Follow Up Closed - In	No
CASTRATE-RESISTANT PROSTATE CANCER (MAINSAIL) A Phase 3, Multicentre, Randomised, Open-Label, Parallel-Group Study of the Efficacy and Safety of Lenalidomide (Revlimid) Versus Chlorambucil as First-Line Therapy for Previously Untreated Elderly Patients with B-Cell Chronic Lymphocytic Leukaemia. (The Origin Trial) A Phase 3, Randomised, Double-blind, Placebo-controlled, Parallel-group, Multi-centre Study to Evaluate the Safety and Efficacy of Ustekinumab Induction Therapy in Particiapnts with Moderately to Severely Active Crohn?s Disease Who Have Failed or Are Intolerant to TNF Antagonist Therapy A Phase 3, Randomized, Double-Blind Study to Evaluate the Safety and	09/H1107/103	15 Sponsor Follow Up Closed - In 5 31/05/2011 Follow Up Closed - In	No
CASTRATE-RESISTANT PROSTATE CANCER (MAINSAIL) A Phase 3, Multicentre, Randomised, Open-Label, Parallel-Group Study of the Efficacy and Safety of Lenalidomide (Revlimid) Versus Chlorambucil as First-Line Therapy for Previously Untreated Elderly Patients with B-Cell Chronic Lymphocytic Leukaemia. (The Origin Trial) A Phase 3, Randomised, Double-blind, Placebo-controlled, Parallel-group, Multi-centre Study to Evaluate the Safety and Efficacy of Ustekinumab Induction Therapy in Particiapnts with Moderately to Severely Active Crohn?s Disease Who Have Failed or Are Intolerant to TNF Antagonist Therapy A Phase 3, Randomized, Double-Blind Study to Evaluate the Safety and Efficacy of GS 9350-boosted Atazanavir Versus Ritonavir-boosted Atazanavir Each Administered with Emtricitabine/Tenofovir Disoproxil Fumarate in HIV 1	09/H1107/103 11/SC/0326	15 Sponsor Follow Up Closed - In 5 31/05/2011 Follow Up Closed - In 5 15/09/2013 Follow Up Closed - In	No
CASTRATE-RESISTANT PROSTATE CANCER (MAINSAIL) A Phase 3, Multicentre, Randomised, Open-Label, Parallel-Group Study of the Efficacy and Safety of Lenalidomide (Revlimid) Versus Chlorambucil as First-Line Therapy for Previously Untreated Elderly Patients with B-Cell Chronic Lymphocytic Leukaemia. (The Origin Trial) A Phase 3, Randomised, Double-blind, Placebo-controlled, Parallel-group, Multi-centre Study to Evaluate the Safety and Efficacy of Ustekinumab Induction Therapy in Particiapnts with Moderately to Severely Active Crohn?s Disease Who Have Failed or Are Intolerant to TNF Antagonist Therapy A Phase 3, Randomized, Double-Blind Study to Evaluate the Safety and Efficacy of GS 9350-boosted Atazanavir Versus Ritonavir-boosted Atazanavir Each Administered with Emtricitabine/Tenofovir Disoproxil Fumarate in HIV 1 Infected, Antiretroviral Treatment-Naïve Adults. (GS-US-216-0114)	09/H1107/103	15 Sponsor Follow Up Closed - In 5 31/05/2011 Follow Up Closed - In 5 15/09/2013 Follow Up Closed - In	No No
CASTRATE-RESISTANT PROSTATE CANCER (MAINSAIL) A Phase 3, Multicentre, Randomised, Open-Label, Parallel-Group Study of the Efficacy and Safety of Lenalidomide (Revlimid) Versus Chlorambucil as First-Line Therapy for Previously Untreated Elderly Patients with B-Cell Chronic Lymphocytic Leukaemia. (The Origin Trial) A Phase 3, Randomised, Double-blind, Placebo-controlled, Parallel-group, Multi-centre Study to Evaluate the Safety and Efficacy of Ustekinumab Induction Therapy in Particiapnts with Moderately to Severely Active Crohn?s Disease Who Have Failed or Are Intolerant to TNF Antagonist Therapy A Phase 3, Randomized, Double-Blind Study to Evaluate the Safety and Efficacy of GS 9350-boosted Atazanavir Versus Ritonavir-boosted Atazanavir Each Administered with Emtricitabine/Tenofovir Disoproxil Fumarate in HIV 1 Infected, Antiretroviral Treatment-Naïve Adults. (GS-US-216-0114) A Phase III Prospective, Two-cohort, Non-randomized, Multi-centre, Multi-	09/H1107/103 11/SC/0326	15 Sponsor Follow Up Closed - In 5 31/05/2011 Follow Up Closed - In 5 15/09/2013 Follow Up Closed - In 8 04/06/2010 Follow Up	No No
CASTRATE-RESISTANT PROSTATE CANCER (MAINSAIL) A Phase 3, Multicentre, Randomised, Open-Label, Parallel-Group Study of the Efficacy and Safety of Lenalidomide (Revlimid) Versus Chlorambucil as First-Line Therapy for Previously Untreated Elderly Patients with B-Cell Chronic Lymphocytic Leukaemia. (The Origin Trial) A Phase 3, Randomised, Double-blind, Placebo-controlled, Parallel-group, Multi-centre Study to Evaluate the Safety and Efficacy of Ustekinumab Induction Therapy in Particiapnts with Moderately to Severely Active Crohn?s Disease Who Have Failed or Are Intolerant to TNF Antagonist Therapy A Phase 3, Randomized, Double-Blind Study to Evaluate the Safety and Efficacy of GS 9350-boosted Atazanavir Versus Ritonavir-boosted Atazanavir Each Administered with Emtricitabine/Tenofovir Disoproxil Fumarate in HIV 1 Infected, Antiretroviral Treatment-Naïve Adults. (GS-US-216-0114) A Phase III Prospective, Two-cohort, Non-randomized, Multi-centre, Multinational, Open Label Study to Assess the Safety of Assisted- and Self-	09/H1107/103 11/SC/0326	15 Sponsor Follow Up Closed - In 5 31/05/2011 Follow Up Closed - In 5 15/09/2013 Follow Up Closed - In 8 04/06/2010 Follow Up No date	No No
CASTRATE-RESISTANT PROSTATE CANCER (MAINSAIL) A Phase 3, Multicentre, Randomised, Open-Label, Parallel-Group Study of the Efficacy and Safety of Lenalidomide (Revlimid) Versus Chlorambucil as First-Line Therapy for Previously Untreated Elderly Patients with B-Cell Chronic Lymphocytic Leukaemia. (The Origin Trial) A Phase 3, Randomised, Double-blind, Placebo-controlled, Parallel-group, Multi-centre Study to Evaluate the Safety and Efficacy of Ustekinumab Induction Therapy in Particiapnts with Moderately to Severely Active Crohn?s Disease Who Have Failed or Are Intolerant to TNF Antagonist Therapy A Phase 3, Randomized, Double-Blind Study to Evaluate the Safety and Efficacy of GS 9350-boosted Atazanavir Versus Ritonavir-boosted Atazanavir Each Administered with Emtricitabine/Tenofovir Disoproxil Fumarate in HIV 1 Infected, Antiretroviral Treatment-Naïve Adults. (GS-US-216-0114) A Phase III Prospective, Two-cohort, Non-randomized, Multi-centre, Multinational, Open Label Study to Assess the Safety of Assisted- and Selfadministered Subcutaneous Trastuzumab as Adjuvant Therapy in Patients	09/H1107/103 11/SC/0326 10/H0711/34	Closed - In 5 31/05/2011 Follow Up Closed - In 5 15/09/2013 Follow Up Closed - In 6 04/06/2010 Follow Up No date agreed with Closed - In	No No
CASTRATE-RESISTANT PROSTATE CANCER (MAINSAIL) A Phase 3, Multicentre, Randomised, Open-Label, Parallel-Group Study of the Efficacy and Safety of Lenalidomide (Revlimid) Versus Chlorambucil as First-Line Therapy for Previously Untreated Elderly Patients with B-Cell Chronic Lymphocytic Leukaemia. (The Origin Trial) A Phase 3, Randomised, Double-blind, Placebo-controlled, Parallel-group, Multi-centre Study to Evaluate the Safety and Efficacy of Ustekinumab Induction Therapy in Particiapnts with Moderately to Severely Active Crohn?s Disease Who Have Failed or Are Intolerant to TNF Antagonist Therapy A Phase 3, Randomized, Double-Blind Study to Evaluate the Safety and Efficacy of GS 9350-boosted Atazanavir Versus Ritonavir-boosted Atazanavir Each Administered with Emtricitabine/Tenofovir Disoproxil Fumarate in HIV 1 Infected, Antiretroviral Treatment-Naïve Adults. (GS-US-216-0114) A Phase III Prospective, Two-cohort, Non-randomized, Multi-centre, Multinational, Open Label Study to Assess the Safety of Assisted- and Self-	09/H1107/103 11/SC/0326	15 Sponsor Follow Up Closed - In 5 31/05/2011 Follow Up Closed - In 5 15/09/2013 Follow Up Closed - In 8 04/06/2010 Follow Up No date	No No
CASTRATE-RESISTANT PROSTATE CANCER (MAINSAIL) A Phase 3, Multicentre, Randomised, Open-Label, Parallel-Group Study of the Efficacy and Safety of Lenalidomide (Revlimid) Versus Chlorambucil as First-Line Therapy for Previously Untreated Elderly Patients with B-Cell Chronic Lymphocytic Leukaemia. (The Origin Trial) A Phase 3, Randomised, Double-blind, Placebo-controlled, Parallel-group, Multi-centre Study to Evaluate the Safety and Efficacy of Ustekinumab Induction Therapy in Particiapnts with Moderately to Severely Active Crohn?s Disease Who Have Failed or Are Intolerant to TNF Antagonist Therapy A Phase 3, Randomized, Double-Blind Study to Evaluate the Safety and Efficacy of GS 9350-boosted Atazanavir Versus Ritonavir-boosted Atazanavir Each Administered with Emtricitabine/Tenofovir Disoproxil Fumarate in HIV 1 Infected, Antiretroviral Treatment-Naïve Adults. (GS-US-216-0114) A Phase III Prospective, Two-cohort, Non-randomized, Multi-centre, Multinational, Open Label Study to Assess the Safety of Assisted- and Selfadministered Subcutaneous Trastuzumab as Adjuvant Therapy in Patients with Operable HER2-positive Early Breast Cancer	09/H1107/103 11/SC/0326 10/H0711/34	Closed - In 5 31/05/2011 Follow Up Closed - In 5 15/09/2013 Follow Up Closed - In 6 04/06/2010 Follow Up No date agreed with Closed - In	No No
CASTRATE-RESISTANT PROSTATE CANCER (MAINSAIL) A Phase 3, Multicentre, Randomised, Open-Label, Parallel-Group Study of the Efficacy and Safety of Lenalidomide (Revlimid) Versus Chlorambucil as First-Line Therapy for Previously Untreated Elderly Patients with B-Cell Chronic Lymphocytic Leukaemia. (The Origin Trial) A Phase 3, Randomised, Double-blind, Placebo-controlled, Parallel-group, Multi-centre Study to Evaluate the Safety and Efficacy of Ustekinumab Induction Therapy in Particiapnts with Moderately to Severely Active Crohn?s Disease Who Have Failed or Are Intolerant to TNF Antagonist Therapy A Phase 3, Randomized, Double-Blind Study to Evaluate the Safety and Efficacy of GS 9350-boosted Atazanavir Versus Ritonavir-boosted Atazanavir Each Administered with Emtricitabine/Tenofovir Disoproxil Fumarate in HIV 1 Infected, Antiretroviral Treatment-Naïve Adults. (GS-US-216-0114) A Phase III Prospective, Two-cohort, Non-randomized, Multi-centre, Multinational, Open Label Study to Assess the Safety of Assisted- and Selfadministered Subcutaneous Trastuzumab as Adjuvant Therapy in Patients with Operable HER2-positive Early Breast Cancer	09/H1107/103 11/SC/0326 10/H0711/34	Closed - In 5 31/05/2011 Follow Up Closed - In 5 15/09/2013 Follow Up Closed - In 6 04/06/2010 Follow Up No date agreed with Closed - In	No No
CASTRATE-RESISTANT PROSTATE CANCER (MAINSAIL) A Phase 3, Multicentre, Randomised, Open-Label, Parallel-Group Study of the Efficacy and Safety of Lenalidomide (Revlimid) Versus Chlorambucil as First-Line Therapy for Previously Untreated Elderly Patients with B-Cell Chronic Lymphocytic Leukaemia. (The Origin Trial) A Phase 3, Randomised, Double-blind, Placebo-controlled, Parallel-group, Multi-centre Study to Evaluate the Safety and Efficacy of Ustekinumab Induction Therapy in Particiapnts with Moderately to Severely Active Crohn?s Disease Who Have Failed or Are Intolerant to TNF Antagonist Therapy A Phase 3, Randomized, Double-Blind Study to Evaluate the Safety and Efficacy of GS 9350-boosted Atazanavir Versus Ritonavir-boosted Atazanavir Each Administered with Emtricitabine/Tenofovir Disoproxil Fumarate in HIV 1 Infected, Antiretroviral Treatment-Naïve Adults. (GS-US-216-0114) A Phase III Prospective, Two-cohort, Non-randomized, Multi-centre, Multinational, Open Label Study to Assess the Safety of Assisted- and Selfadministered Subcutaneous Trastuzumab as Adjuvant Therapy in Patients with Operable HER2-positive Early Breast Cancer A PHASE III, RANDOMIZED, DOUBLE-BLIND, PLACEBO CONTROLLED CLINICAL TRIAL TO EVALUATE THE EFFICACY AND SAFETY OF PERTUZUMAB +	09/H1107/103 11/SC/0326 10/H0711/34	Closed - In 5 31/05/2011 Follow Up Closed - In 5 15/09/2013 Follow Up Closed - In 8 04/06/2010 Follow Up No date agreed with Closed - In 6 Sponsor Follow Up	No No
CASTRATE-RESISTANT PROSTATE CANCER (MAINSAIL) A Phase 3, Multicentre, Randomised, Open-Label, Parallel-Group Study of the Efficacy and Safety of Lenalidomide (Revlimid) Versus Chlorambucil as First-Line Therapy for Previously Untreated Elderly Patients with B-Cell Chronic Lymphocytic Leukaemia. (The Origin Trial) A Phase 3, Randomised, Double-blind, Placebo-controlled, Parallel-group, Multi-centre Study to Evaluate the Safety and Efficacy of Ustekinumab Induction Therapy in Particiapnts with Moderately to Severely Active Crohn?s Disease Who Have Failed or Are Intolerant to TNF Antagonist Therapy A Phase 3, Randomized, Double-Blind Study to Evaluate the Safety and Efficacy of GS 9350-boosted Atazanavir Versus Ritonavir-boosted Atazanavir Each Administered with Emtricitabine/Tenofovir Disoproxil Fumarate in HIV 1 Infected, Antiretroviral Treatment-Naïve Adults. (GS-US-216-0114) A Phase III Prospective, Two-cohort, Non-randomized, Multi-centre, Multinational, Open Label Study to Assess the Safety of Assisted- and Selfadministered Subcutaneous Trastuzumab as Adjuvant Therapy in Patients with Operable HER2-positive Early Breast Cancer	09/H1107/103 11/SC/0326 10/H0711/34	Closed - In 5 31/05/2011 Follow Up Closed - In 5 15/09/2013 Follow Up Closed - In 6 04/06/2010 Follow Up No date agreed with Closed - In	No No
CASTRATE-RESISTANT PROSTATE CANCER (MAINSAIL) A Phase 3, Multicentre, Randomised, Open-Label, Parallel-Group Study of the Efficacy and Safety of Lenalidomide (Revlimid) Versus Chlorambucil as First-Line Therapy for Previously Untreated Elderly Patients with B-Cell Chronic Lymphocytic Leukaemia. (The Origin Trial) A Phase 3, Randomised, Double-blind, Placebo-controlled, Parallel-group, Multi-centre Study to Evaluate the Safety and Efficacy of Ustekinumab Induction Therapy in Particiapnts with Moderately to Severely Active Crohn?s Disease Who Have Failed or Are Intolerant to TNF Antagonist Therapy A Phase 3, Randomized, Double-Blind Study to Evaluate the Safety and Efficacy of GS 9350-boosted Atazanavir Versus Ritonavir-boosted Atazanavir Each Administered with Emtricitabine/Tenofovir Disoproxil Fumarate in HIV 1 Infected, Antiretroviral Treatment-Naïve Adults. (GS-US-216-0114) A Phase III Prospective, Two-cohort, Non-randomized, Multi-centre, Multinational, Open Label Study to Assess the Safety of Assisted- and Selfadministered Subcutaneous Trastuzumab as Adjuvant Therapy in Patients with Operable HER2-positive Early Breast Cancer A PHASE III, RANDOMIZED, DOUBLE-BLIND, PLACEBO CONTROLLED CLINICAL TRIAL TO EVALUATE THE EFFICACY AND SAFETY OF PERTUZUMAB +	09/H1107/103 11/SC/0326 10/H0711/34	Closed - In 5 31/05/2011 Follow Up Closed - In 5 15/09/2013 Follow Up Closed - In 8 04/06/2010 Follow Up No date agreed with Closed - In 6 Sponsor Follow Up	No No
CASTRATE-RESISTANT PROSTATE CANCER (MAINSAIL) A Phase 3, Multicentre, Randomised, Open-Label, Parallel-Group Study of the Efficacy and Safety of Lenalidomide (Revlimid) Versus Chlorambucil as First-Line Therapy for Previously Untreated Elderly Patients with B-Cell Chronic Lymphocytic Leukaemia. (The Origin Trial) A Phase 3, Randomised, Double-blind, Placebo-controlled, Parallel-group, Multi-centre Study to Evaluate the Safety and Efficacy of Ustekinumab Induction Therapy in Particiapnts with Moderately to Severely Active Crohn?s Disease Who Have Failed or Are Intolerant to TNF Antagonist Therapy A Phase 3, Randomized, Double-Blind Study to Evaluate the Safety and Efficacy of GS 9350-boosted Atazanavir Versus Ritonavir-boosted Atazanavir Each Administered with Emtricitabine/Tenofovir Disoproxil Fumarate in HIV 1 Infected, Antiretroviral Treatment-Naïve Adults. (GS-US-216-0114) A Phase III Prospective, Two-cohort, Non-randomized, Multi-centre, Multinational, Open Label Study to Assess the Safety of Assisted- and Selfadministered Subcutaneous Trastuzumab as Adjuvant Therapy in Patients with Operable HER2-positive Early Breast Cancer A PHASE III, RANDOMIZED, DOUBLE-BLIND, PLACEBO CONTROLLED CLINICAL TRIAL TO EVALUATE THE EFFICACY AND SAFETY OF PERTUZUMAB + TRASTUZUMAB + DOCETAXEL IN	09/H1107/103 11/SC/0326 10/H0711/34 12/SC/0139	Closed - In 5 31/05/2011 Follow Up Closed - In 5 15/09/2013 Follow Up Closed - In 8 04/06/2010 Follow Up No date agreed with Closed - In 6 Sponsor Follow Up Closed - In	No No
CASTRATE-RESISTANT PROSTATE CANCER (MAINSAIL) A Phase 3, Multicentre, Randomised, Open-Label, Parallel-Group Study of the Efficacy and Safety of Lenalidomide (Revlimid) Versus Chlorambucil as First-Line Therapy for Previously Untreated Elderly Patients with B-Cell Chronic Lymphocytic Leukaemia. (The Origin Trial) A Phase 3, Randomised, Double-blind, Placebo-controlled, Parallel-group, Multi-centre Study to Evaluate the Safety and Efficacy of Ustekinumab Induction Therapy in Particiapnts with Moderately to Severely Active Crohn?s Disease Who Have Failed or Are Intolerant to TNF Antagonist Therapy A Phase 3, Randomized, Double-Blind Study to Evaluate the Safety and Efficacy of GS 9350-boosted Atazanavir Versus Ritonavir-boosted Atazanavir Each Administered with Emtricitabine/Tenofovir Disoproxil Fumarate in HIV 1 Infected, Antiretroviral Treatment-Naïve Adults. (GS-US-216-0114) A Phase III Prospective, Two-cohort, Non-randomized, Multi-centre, Multinational, Open Label Study to Assess the Safety of Assisted- and Selfadministered Subcutaneous Trastuzumab as Adjuvant Therapy in Patients with Operable HER2-positive Early Breast Cancer A PHASE III, RANDOMIZED, DOUBLE-BLIND, PLACEBO CONTROLLED CLINICAL TRIAL TO EVALUATE THE EFFICACY AND SAFETY OF PERTUZUMAB + DOCETAXEL IN	09/H1107/103 11/SC/0326 10/H0711/34 12/SC/0139	Closed - In 5 31/05/2011 Follow Up Closed - In 5 15/09/2013 Follow Up Closed - In 8 04/06/2010 Follow Up No date agreed with Closed - In 6 Sponsor Follow Up Closed - In	No No
CASTRATE-RESISTANT PROSTATE CANCER (MAINSAIL) A Phase 3, Multicentre, Randomised, Open-Label, Parallel-Group Study of the Efficacy and Safety of Lenalidomide (Revlimid) Versus Chlorambucil as First-Line Therapy for Previously Untreated Elderly Patients with B-Cell Chronic Lymphocytic Leukaemia. (The Origin Trial) A Phase 3, Randomised, Double-blind, Placebo-controlled, Parallel-group, Multi-centre Study to Evaluate the Safety and Efficacy of Ustekinumab Induction Therapy in Particiapnts with Moderately to Severely Active Crohn?s Disease Who Have Failed or Are Intolerant to TNF Antagonist Therapy A Phase 3, Randomized, Double-Blind Study to Evaluate the Safety and Efficacy of GS 9350-boosted Atazanavir Versus Ritonavir-boosted Atazanavir Each Administered with Emtricitabine/Tenofovir Disoproxil Fumarate in HIV 1 Infected, Antiretroviral Treatment-Naïve Adults. (GS-US-216-0114) A Phase III Prospective, Two-cohort, Non-randomized, Multi-centre, Multinational, Open Label Study to Assess the Safety of Assisted- and Selfadministered Subcutaneous Trastuzumab as Adjuvant Therapy in Patients with Operable HER2-positive Early Breast Cancer A PHASE III, RANDOMIZED, DOUBLE-BLIND, PLACEBO CONTROLLED CLINICAL TRIAL TO EVALUATE THE EFFICACY AND SAFETY OF PERTUZUMAB + TRASTUZUMAB + DOCETAXEL IN PREVIOUSLY UNTREATED HER2-POSITIVE METASTATIC BREAST CANCER	09/H1107/103 11/SC/0326 10/H0711/34 12/SC/0139	Closed - In 5 31/05/2011 Follow Up Closed - In 5 15/09/2013 Follow Up Closed - In 8 04/06/2010 Follow Up No date agreed with Closed - In 6 Sponsor Follow Up Closed - In 7 Sollow Up Closed - In 7 Sollow Up	No No
CASTRATE-RESISTANT PROSTATE CANCER (MAINSAIL) A Phase 3, Multicentre, Randomised, Open-Label, Parallel-Group Study of the Efficacy and Safety of Lenalidomide (Revlimid) Versus Chlorambucil as First-Line Therapy for Previously Untreated Elderly Patients with B-Cell Chronic Lymphocytic Leukaemia. (The Origin Trial) A Phase 3, Randomised, Double-blind, Placebo-controlled, Parallel-group, Multi-centre Study to Evaluate the Safety and Efficacy of Ustekinumab Induction Therapy in Particiapnts with Moderately to Severely Active Crohn?s Disease Who Have Failed or Are Intolerant to TNF Antagonist Therapy A Phase 3, Randomized, Double-Blind Study to Evaluate the Safety and Efficacy of GS 9350-boosted Atazanavir Versus Ritonavir-boosted Atazanavir Each Administered with Emtricitabine/Tenofovir Disoproxil Fumarate in HIV 1 Infected, Antiretroviral Treatment-Naïve Adults. (GS-US-216-0114) A Phase III Prospective, Two-cohort, Non-randomized, Multi-centre, Multinational, Open Label Study to Assess the Safety of Assisted- and Self-administered Subcutaneous Trastuzumab as Adjuvant Therapy in Patients with Operable HER2-positive Early Breast Cancer A PHASE III, RANDOMIZED, DOUBLE-BLIND, PLACEBO CONTROLLED CLINICAL TRIAL TO EVALUATE THE EFFICACY AND SAFETY OF PERTUZUMAB + TRASTUZUMAB + DOCETAXEL IN PREVIOUSLY UNTREATED HER2-POSITIVE METASTATIC BREAST CANCER	09/H1107/103 11/SC/0326 10/H0711/34 12/SC/0139	Closed - In 5 31/05/2011 Follow Up Closed - In 5 15/09/2013 Follow Up Closed - In 8 04/06/2010 Follow Up No date agreed with Closed - In 6 Sponsor Follow Up Closed - In 7 Closed - In 7 Closed - In 8 O4/06/2010 Follow Up	No No

A Bandamized Multicenter Phase 2 Study Comparing Carfillomia		N	lo data		
A Randomized, Multicenter, Phase 3 Study Comparing Carfilzomib,			lo date	Closed - In	
Lenalidomide, and Dexamethasone (CRd) vs Lenalidomide and Dexamethasone (Rd) in Subjects with Relapsed Multiple Myeloma	10/H0806/71		greed with ponsor	Follow Up	No
Dexamethasone (kd) in Subjects with kelapsed Multiple Myeloma	10/110800/71	<i>J</i> 3	porisor	rollow op	INU
A Randomized, Open-label, Phase 3 Study of Carfilzomib vs Best Supportive				Closed - In	
Care in Subjects with Relapsed and Refractory Multiple Myeloma	11/SC/0479	4	30/11/2012		No
	11/30/04/3	4	30/11/2012	TOHOW OP	INU
An Open-Label, Multicenter, Follow-up Study to Evaluate the Long-Term				0	
Safety and Efficacy of Brivaracetam Used as Adjunctive Treatment in Subjects	44 /NE /0000	_	20/44/2042	Closed - In	
Aged 16 Years or Older with Partial Onset Seizures	11/NE/0039	5	30/11/2012	Follow Up	No
An open-label, randomized, multi-center, Phase III study to compare the					
safety and efficacy of TKI258 versus sorafenib in patients with metastatic					
renal cell carcinoma after failure of antiangiogenic (VEGF-targeted and mTOR				Closed - In	
inhibitor) therapies (GOLD)	11/LO/0548	4	31/10/2012	Follow Up	No
CIS Study: An international, Multi-centre, Randomised, Double-Blind, placebo					
controlled parallel group study to evaluate the efficacy and safety of two year		N	lo date		
treatment with Teriflunomide 7mg once daily and 14mg once daily versus the		a	greed with	Closed - In	
plaebo in patients with a first clinical episode suggestive of multiple sclerosis	07/H0707/112	7 S	ponsor	Follow Up	No
		N	lo date		
CoreValve Advance International Post Market Study A Multi-centre,		a	greed with	Closed - In	
Interventional, Prospective, Post-Market Release Study	10/H0724/33	100 S	ponsor	Follow Up	No
Phase II randomised, placebo controlled trial to investigate repeat dose		N	lo date		
antimicrobial photodynamic therapy in patients with chronic venous leg		a	greed with	Closed - In	
ulcers	08/H0206/68	5 S	ponsor	Follow Up	No
TEMSO Extension: Long-term extension study of the multinational, double-					
blind, placebo controlled study EFC6049 (HMR1726D/3001) to document the		N	lo date		
safety of two doses of teriflunomide (7 and 14mg) in patients with multiple		a	greed with	Closed - In	
sclerosis with relapses.	06/MRE04/84	10 S	ponsor	Follow Up	No
A Phase III Randomized, Double-blind Study of the Safety and Efficacy of					
GSK1349572 50 mg Once Daily Versus Raltegravir 400 mg Twice Daily, Both					
Administered with an Investigator-selected Background Regimen Over 48					
Weeks in HIV-1 Infected, Integrase Inhibitor-Naïve, Antiretroviral Therapy-				Closed - Follow	
Experienced Adults ING111762	10/H0706/67	1	07/05/2015	Up Complete	Yes
A Double-Blind, Randomized, Parallel Group, Multi-Centre Study to Evaluate	10/110700/07		07/03/2013	op complete	103
the Efficacy and Safety of Mirabegron Compared to Solifenacin in Subjects					
with Overactive Bladder (OAB) Treated with Antimuscarinics and Dissatisfied				Closed - Follow	
due to Lack of Efficacy	12/LO/0410	6	30/11/2012	Up Complete	No
due to Eack of Efficacy	12/10/0410	- 0	30/11/2012	op complete	INO
A multicentre, single arm study of Trastuzumab Emtasine (TDM-1) in HER2					
				Closed Follow	
positive locally advanced or metastatic breast cancer patients who have	12/10/1251	-	20/00/2012	Closed - Follow	Na
received prior anti-HER2 and chemotherapy-based treatment.	12/LO/1351	5	30/09/2013	Closed - Follow Up Complete	No
received prior anti-HER2 and chemotherapy-based treatment.	12/LO/1351	5	30/09/2013	Up Complete	No
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration				Up Complete Closed - Follow	
received prior anti-HER2 and chemotherapy-based treatment.	12/LO/1351 10/H1107/42	5		Up Complete	No No
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration Of QUTENZA For The Treatment Of Peripheral Neuropathic Pain				Up Complete Closed - Follow	
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration Of QUTENZA For The Treatment Of Peripheral Neuropathic Pain A Phase 3, Randomized, Double-Blind, Double-Dummy, Parallel Group, Multi-				Up Complete Closed - Follow	
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration Of QUTENZA For The Treatment Of Peripheral Neuropathic Pain A Phase 3, Randomized, Double-Blind, Double-Dummy, Parallel Group, Multi-Center, Multi-National Study for Evaluation of Efficacy and Safety of DU-176b				Up Complete Closed - Follow Up Complete	
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration Of QUTENZA For The Treatment Of Peripheral Neuropathic Pain A Phase 3, Randomized, Double-Blind, Double-Dummy, Parallel Group, Multi-Center, Multi-National Study for Evaluation of Efficacy and Safety of DU-176b Versus Warfarin In Subjects with Atrial Fibrillation – Effective Anticoagulation	10/H1107/42	6	31/08/2012	Up Complete Closed - Follow Up Complete Closed - Follow	
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration Of QUTENZA For The Treatment Of Peripheral Neuropathic Pain A Phase 3, Randomized, Double-Blind, Double-Dummy, Parallel Group, Multi-Center, Multi-National Study for Evaluation of Efficacy and Safety of DU-176b			31/08/2012	Up Complete Closed - Follow Up Complete	
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration Of QUTENZA For The Treatment Of Peripheral Neuropathic Pain A Phase 3, Randomized, Double-Blind, Double-Dummy, Parallel Group, Multi-Center, Multi-National Study for Evaluation of Efficacy and Safety of DU-176b Versus Warfarin In Subjects with Atrial Fibrillation – Effective Anticoagulation with Factor Xa Next Generation in Atrial Fibrillation (ENGAGE – AF TIMI - 48)	10/H1107/42	6	31/08/2012	Up Complete Closed - Follow Up Complete Closed - Follow	No
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration Of QUTENZA For The Treatment Of Peripheral Neuropathic Pain A Phase 3, Randomized, Double-Blind, Double-Dummy, Parallel Group, Multi-Center, Multi-National Study for Evaluation of Efficacy and Safety of DU-176b Versus Warfarin In Subjects with Atrial Fibrillation – Effective Anticoagulation with Factor Xa Next Generation in Atrial Fibrillation (ENGAGE – AF TIMI - 48) A Randomized, Double-Blind, Placebo-Controlled, Multicenter, Parallel-Group	10/H1107/42	6	31/08/2012	Up Complete Closed - Follow Up Complete Closed - Follow Up Complete	No
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration Of QUTENZA For The Treatment Of Peripheral Neuropathic Pain A Phase 3, Randomized, Double-Blind, Double-Dummy, Parallel Group, Multi-Center, Multi-National Study for Evaluation of Efficacy and Safety of DU-176b Versus Warfarin In Subjects with Atrial Fibrillation – Effective Anticoagulation with Factor Xa Next Generation in Atrial Fibrillation (ENGAGE – AF TIMI - 48) A Randomized, Double-Blind, Placebo-Controlled, Multicenter, Parallel-Group Study to Evaluate the Efficacy and Safety of Brivaracetam in Subjects (=16 To	10/H1107/42 09/H0706/8	10	31/08/2012 18/05/2012	Up Complete Closed - Follow Up Complete Closed - Follow Up Complete Closed - Follow	No No
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration Of QUTENZA For The Treatment Of Peripheral Neuropathic Pain A Phase 3, Randomized, Double-Blind, Double-Dummy, Parallel Group, Multi-Center, Multi-National Study for Evaluation of Efficacy and Safety of DU-176b Versus Warfarin In Subjects with Atrial Fibrillation – Effective Anticoagulation with Factor Xa Next Generation in Atrial Fibrillation (ENGAGE – AF TIMI - 48) A Randomized, Double-Blind, Placebo-Controlled, Multicenter, Parallel-Group Study to Evaluate the Efficacy and Safety of Brivaracetam in Subjects (=16 To 80 Years Old) With Partial Onset Seizures	10/H1107/42	6 10 5	31/08/2012 18/05/2012 02/05/2012	Up Complete Closed - Follow Up Complete Closed - Follow Up Complete	No
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration Of QUTENZA For The Treatment Of Peripheral Neuropathic Pain A Phase 3, Randomized, Double-Blind, Double-Dummy, Parallel Group, Multi-Center, Multi-National Study for Evaluation of Efficacy and Safety of DU-176b Versus Warfarin In Subjects with Atrial Fibrillation – Effective Anticoagulation with Factor Xa Next Generation in Atrial Fibrillation (ENGAGE – AF TIMI - 48) A Randomized, Double-Blind, Placebo-Controlled, Multicenter, Parallel-Group Study to Evaluate the Efficacy and Safety of Brivaracetam in Subjects (=16 To 80 Years Old) With Partial Onset Seizures A single arm, open-label, multicenter study evaluating the long-term safety	10/H1107/42 09/H0706/8	6 10 5	31/08/2012 18/05/2012	Up Complete Closed - Follow Up Complete Closed - Follow Up Complete Closed - Follow Up Complete	No No
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration Of QUTENZA For The Treatment Of Peripheral Neuropathic Pain A Phase 3, Randomized, Double-Blind, Double-Dummy, Parallel Group, Multi-Center, Multi-National Study for Evaluation of Efficacy and Safety of DU-176b Versus Warfarin In Subjects with Atrial Fibrillation – Effective Anticoagulation with Factor Xa Next Generation in Atrial Fibrillation (ENGAGE – AF TIMI - 48) A Randomized, Double-Blind, Placebo-Controlled, Multicenter, Parallel-Group Study to Evaluate the Efficacy and Safety of Brivaracetam in Subjects (=16 To 80 Years Old) With Partial Onset Seizures	10/H1107/42 09/H0706/8	6 10 5 N	31/08/2012 18/05/2012 02/05/2012	Up Complete Closed - Follow Up Complete Closed - Follow Up Complete Closed - Follow	No No
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration Of QUTENZA For The Treatment Of Peripheral Neuropathic Pain A Phase 3, Randomized, Double-Blind, Double-Dummy, Parallel Group, Multi-Center, Multi-National Study for Evaluation of Efficacy and Safety of DU-176b Versus Warfarin In Subjects with Atrial Fibrillation – Effective Anticoagulation with Factor Xa Next Generation in Atrial Fibrillation (ENGAGE – AF TIMI - 48) A Randomized, Double-Blind, Placebo-Controlled, Multicenter, Parallel-Group Study to Evaluate the Efficacy and Safety of Brivaracetam in Subjects (=16 To 80 Years Old) With Partial Onset Seizures A single arm, open-label, multicenter study evaluating the long-term safety	10/H1107/42 09/H0706/8	10 5 N	31/08/2012 18/05/2012 02/05/2012 lo date	Up Complete Closed - Follow Up Complete Closed - Follow Up Complete Closed - Follow Up Complete	No No
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration Of QUTENZA For The Treatment Of Peripheral Neuropathic Pain A Phase 3, Randomized, Double-Blind, Double-Dummy, Parallel Group, Multi-Center, Multi-National Study for Evaluation of Efficacy and Safety of DU-176b Versus Warfarin In Subjects with Atrial Fibrillation – Effective Anticoagulation with Factor Xa Next Generation in Atrial Fibrillation (ENGAGE – AF TIMI - 48) A Randomized, Double-Blind, Placebo-Controlled, Multicenter, Parallel-Group Study to Evaluate the Efficacy and Safety of Brivaracetam in Subjects (=16 To 80 Years Old) With Partial Onset Seizures A single arm, open-label, multicenter study evaluating the long-term safety and tolerability of 0.5 mg fingolimod (FTY720) administered orally once daily	10/H1107/42 09/H0706/8 11/NE/0038	10 5 N	31/08/2012 18/05/2012 02/05/2012 lo date greed with	Up Complete Closed - Follow Up Complete	No No
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration Of QUTENZA For The Treatment Of Peripheral Neuropathic Pain A Phase 3, Randomized, Double-Blind, Double-Dummy, Parallel Group, Multi-Center, Multi-National Study for Evaluation of Efficacy and Safety of DU-176b Versus Warfarin In Subjects with Atrial Fibrillation – Effective Anticoagulation with Factor Xa Next Generation in Atrial Fibrillation (ENGAGE – AF TIMI - 48) A Randomized, Double-Blind, Placebo-Controlled, Multicenter, Parallel-Group Study to Evaluate the Efficacy and Safety of Brivaracetam in Subjects (=16 To 80 Years Old) With Partial Onset Seizures A single arm, open-label, multicenter study evaluating the long-term safety and tolerability of 0.5 mg fingolimod (FTY720) administered orally once daily	10/H1107/42 09/H0706/8 11/NE/0038	10 5 N	31/08/2012 18/05/2012 02/05/2012 lo date greed with	Up Complete Closed - Follow Up Complete	No No
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration Of QUTENZA For The Treatment Of Peripheral Neuropathic Pain A Phase 3, Randomized, Double-Blind, Double-Dummy, Parallel Group, Multi-Center, Multi-National Study for Evaluation of Efficacy and Safety of DU-176b Versus Warfarin In Subjects with Atrial Fibrillation – Effective Anticoagulation with Factor Xa Next Generation in Atrial Fibrillation (ENGAGE – AF TIMI - 48) A Randomized, Double-Blind, Placebo-Controlled, Multicenter, Parallel-Group Study to Evaluate the Efficacy and Safety of Brivaracetam in Subjects (=16 To 80 Years Old) With Partial Onset Seizures A single arm, open-label, multicenter study evaluating the long-term safety and tolerability of 0.5 mg fingolimod (FTY720) administered orally once daily in patients with relapsing forms of multiple sclerosis	10/H1107/42 09/H0706/8 11/NE/0038	10 5 N	31/08/2012 18/05/2012 02/05/2012 lo date greed with	Up Complete Closed - Follow Up Complete	No No
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration Of QUTENZA For The Treatment Of Peripheral Neuropathic Pain A Phase 3, Randomized, Double-Blind, Double-Dummy, Parallel Group, Multi-Center, Multi-National Study for Evaluation of Efficacy and Safety of DU-176b Versus Warfarin In Subjects with Atrial Fibrillation – Effective Anticoagulation with Factor Xa Next Generation in Atrial Fibrillation (ENGAGE – AF TIMI - 48) A Randomized, Double-Blind, Placebo-Controlled, Multicenter, Parallel-Group Study to Evaluate the Efficacy and Safety of Brivaracetam in Subjects (=16 To 80 Years Old) With Partial Onset Seizures A single arm, open-label, multicenter study evaluating the long-term safety and tolerability of 0.5 mg fingolimod (FTY720) administered orally once daily in patients with relapsing forms of multiple sclerosis	10/H1107/42 09/H0706/8 11/NE/0038	10 5 N	31/08/2012 18/05/2012 02/05/2012 lo date greed with	Up Complete Closed - Follow Up Complete	No No
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration Of QUTENZA For The Treatment Of Peripheral Neuropathic Pain A Phase 3, Randomized, Double-Blind, Double-Dummy, Parallel Group, Multi-Center, Multi-National Study for Evaluation of Efficacy and Safety of DU-176b Versus Warfarin In Subjects with Atrial Fibrillation – Effective Anticoagulation with Factor Xa Next Generation in Atrial Fibrillation (ENGAGE – AF TIMI - 48) A Randomized, Double-Blind, Placebo-Controlled, Multicenter, Parallel-Group Study to Evaluate the Efficacy and Safety of Brivaracetam in Subjects (=16 To 80 Years Old) With Partial Onset Seizures A single arm, open-label, multicenter study evaluating the long-term safety and tolerability of 0.5 mg fingolimod (FTY720) administered orally once daily in patients with relapsing forms of multiple sclerosis ALTITUDE: A randomised, double-blind, placebo-controlled, parallel-group study to determine whether, in patients with type 2 diabeted at high risk for	10/H1107/42 09/H0706/8 11/NE/0038	10 5 N	31/08/2012 18/05/2012 02/05/2012 lo date greed with ponsor	Up Complete Closed - Follow Up Complete	No No
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration Of QUTENZA For The Treatment Of Peripheral Neuropathic Pain A Phase 3, Randomized, Double-Blind, Double-Dummy, Parallel Group, Multi-Center, Multi-National Study for Evaluation of Efficacy and Safety of DU-176b Versus Warfarin In Subjects with Atrial Fibrillation – Effective Anticoagulation with Factor Xa Next Generation in Atrial Fibrillation (ENGAGE – AF TIMI - 48) A Randomized, Double-Blind, Placebo-Controlled, Multicenter, Parallel-Group Study to Evaluate the Efficacy and Safety of Brivaracetam in Subjects (=16 To 80 Years Old) With Partial Onset Seizures A single arm, open-label, multicenter study evaluating the long-term safety and tolerability of 0.5 mg fingolimod (FTY720) administered orally once daily in patients with relapsing forms of multiple sclerosis ALTITUDE: A randomised, double-blind, placebo-controlled, parallel-group study to determine whether, in patients with type 2 diabeted at high risk for cardiovascular and renal events, aliskiren, on top of conventional treatment,	10/H1107/42 09/H0706/8 11/NE/0038 10/H0904/49	6 10 5 N a 8 S	31/08/2012 18/05/2012 02/05/2012 lo date greed with ponsor	Up Complete Closed - Follow Up Complete	No No No
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration Of QUTENZA For The Treatment Of Peripheral Neuropathic Pain A Phase 3, Randomized, Double-Blind, Double-Dummy, Parallel Group, Multi-Center, Multi-National Study for Evaluation of Efficacy and Safety of DU-176b Versus Warfarin In Subjects with Atrial Fibrillation – Effective Anticoagulation with Factor Xa Next Generation in Atrial Fibrillation (ENGAGE – AF TIMI - 48) A Randomized, Double-Blind, Placebo-Controlled, Multicenter, Parallel-Group Study to Evaluate the Efficacy and Safety of Brivaracetam in Subjects (=16 To 80 Years Old) With Partial Onset Seizures A single arm, open-label, multicenter study evaluating the long-term safety and tolerability of 0.5 mg fingolimod (FTY720) administered orally once daily in patients with relapsing forms of multiple sclerosis ALTITUDE: A randomised, double-blind, placebo-controlled, parallel-group study to determine whether, in patients with type 2 diabeted at high risk for cardiovascular and renal events, aliskiren, on top of conventional treatment,	10/H1107/42 09/H0706/8 11/NE/0038 10/H0904/49	6 10 5 N a 8 S	31/08/2012 18/05/2012 02/05/2012 lo date greed with ponsor	Up Complete Closed - Follow Up Complete	No No No
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration Of QUTENZA For The Treatment Of Peripheral Neuropathic Pain A Phase 3, Randomized, Double-Blind, Double-Dummy, Parallel Group, Multi-Center, Multi-National Study for Evaluation of Efficacy and Safety of DU-176b Versus Warfarin In Subjects with Atrial Fibrillation – Effective Anticoagulation with Factor Xa Next Generation in Atrial Fibrillation (ENGAGE – AF TIMI - 48) A Randomized, Double-Blind, Placebo-Controlled, Multicenter, Parallel-Group Study to Evaluate the Efficacy and Safety of Brivaracetam in Subjects (=16 To 80 Years Old) With Partial Onset Seizures A single arm, open-label, multicenter study evaluating the long-term safety and tolerability of 0.5 mg fingolimod (FTY720) administered orally once daily in patients with relapsing forms of multiple sclerosis ALTITUDE: A randomised, double-blind, placebo-controlled, parallel-group study to determine whether, in patients with type 2 diabeted at high risk for cardiovascular and renal events, aliskiren, on top of conventional treatment, reduces cardiovascular and renal morbidity and mortality	10/H1107/42 09/H0706/8 11/NE/0038 10/H0904/49	6 10 5 N a 8 S	31/08/2012 18/05/2012 02/05/2012 10 date greed with ponsor 01/01/2012	Up Complete Closed - Follow Up Complete	No No No
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration Of QUTENZA For The Treatment Of Peripheral Neuropathic Pain A Phase 3, Randomized, Double-Blind, Double-Dummy, Parallel Group, Multi-Center, Multi-National Study for Evaluation of Efficacy and Safety of DU-176b Versus Warfarin In Subjects with Atrial Fibrillation – Effective Anticoagulation with Factor Xa Next Generation in Atrial Fibrillation (ENGAGE – AF TIMI - 48) A Randomized, Double-Blind, Placebo-Controlled, Multicenter, Parallel-Group Study to Evaluate the Efficacy and Safety of Brivaracetam in Subjects (=16 To 80 Years Old) With Partial Onset Seizures A single arm, open-label, multicenter study evaluating the long-term safety and tolerability of 0.5 mg fingolimod (FTY720) administered orally once daily in patients with relapsing forms of multiple sclerosis ALTITUDE: A randomised, double-blind, placebo-controlled, parallel-group study to determine whether, in patients with type 2 diabeted at high risk for cardiovascular and renal events, aliskiren, on top of conventional treatment, reduces cardiovascular and renal morbidity and mortality FINISH 3: A Phase 3, Randomised, Single Blind, Controlled Trial of Topical Fibrocaps in Intraoperative Surgical Hemostasis	10/H1107/42 09/H0706/8 11/NE/0038 10/H0904/49 07/H0606/126	6 10 5 N a 8 S	31/08/2012 18/05/2012 02/05/2012 10 date greed with ponsor 01/01/2012	Up Complete Closed - Follow Up Complete	No No No
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration Of QUTENZA For The Treatment Of Peripheral Neuropathic Pain A Phase 3, Randomized, Double-Blind, Double-Dummy, Parallel Group, Multi-Center, Multi-National Study for Evaluation of Efficacy and Safety of DU-176b Versus Warfarin In Subjects with Atrial Fibrillation – Effective Anticoagulation with Factor Xa Next Generation in Atrial Fibrillation (ENGAGE – AF TIMI - 48) A Randomized, Double-Blind, Placebo-Controlled, Multicenter, Parallel-Group Study to Evaluate the Efficacy and Safety of Brivaracetam in Subjects (=16 To 80 Years Old) With Partial Onset Seizures A single arm, open-label, multicenter study evaluating the long-term safety and tolerability of 0.5 mg fingolimod (FTY720) administered orally once daily in patients with relapsing forms of multiple sclerosis ALTITUDE: A randomised, double-blind, placebo-controlled, parallel-group study to determine whether, in patients with type 2 diabeted at high risk for cardiovascular and renal events, aliskiren, on top of conventional treatment, reduces cardiovascular and renal morbidity and mortality FINISH 3: A Phase 3, Randomised, Single Blind, Controlled Trial of Topical Fibrocaps in Intraoperative Surgical Hemostasis GWCA0962: A double blind, randomized, placebo-controlled, parallel group	10/H1107/42 09/H0706/8 11/NE/0038 10/H0904/49 07/H0606/126	6 10 5 N a 8 S	31/08/2012 18/05/2012 02/05/2012 10 date greed with ponsor 01/01/2012	Up Complete Closed - Follow Up Complete	No No No
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration Of QUTENZA For The Treatment Of Peripheral Neuropathic Pain A Phase 3, Randomized, Double-Blind, Double-Dummy, Parallel Group, Multi-Center, Multi-National Study for Evaluation of Efficacy and Safety of DU-176b Versus Warfarin In Subjects with Atrial Fibrillation – Effective Anticoagulation with Factor Xa Next Generation in Atrial Fibrillation (ENGAGE – AF TIMI - 48) A Randomized, Double-Blind, Placebo-Controlled, Multicenter, Parallel-Group Study to Evaluate the Efficacy and Safety of Brivaracetam in Subjects (=16 To 80 Years Old) With Partial Onset Seizures A single arm, open-label, multicenter study evaluating the long-term safety and tolerability of 0.5 mg fingolimod (FTY720) administered orally once daily in patients with relapsing forms of multiple sclerosis ALTITUDE: A randomised, double-blind, placebo-controlled, parallel-group study to determine whether, in patients with type 2 diabeted at high risk for cardiovascular and renal events, aliskiren, on top of conventional treatment, reduces cardiovascular and renal morbidity and mortality FINISH 3: A Phase 3, Randomised, Single Blind, Controlled Trial of Topical Fibrocaps in Intraoperative Surgical Hemostasis GWCA0962: A double blind, randomized, placebo-controlled, parallel group study of Sativex oromucosal spray (Sativex®; Nabiximols) as adjunctive	10/H1107/42 09/H0706/8 11/NE/0038 10/H0904/49 07/H0606/126	6 10 5 N a 8 S	31/08/2012 18/05/2012 02/05/2012 10 date greed with ponsor 01/01/2012	Up Complete Closed - Follow Up Complete	No No No
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration Of QUTENZA For The Treatment Of Peripheral Neuropathic Pain A Phase 3, Randomized, Double-Blind, Double-Dummy, Parallel Group, Multi-Center, Multi-National Study for Evaluation of Efficacy and Safety of DU-176b Versus Warfarin In Subjects with Atrial Fibrillation – Effective Anticoagulation with Factor Xa Next Generation in Atrial Fibrillation (ENGAGE – AF TIMI - 48) A Randomized, Double-Blind, Placebo-Controlled, Multicenter, Parallel-Group Study to Evaluate the Efficacy and Safety of Brivaracetam in Subjects (=16 To 80 Years Old) With Partial Onset Seizures A single arm, open-label, multicenter study evaluating the long-term safety and tolerability of 0.5 mg fingolimod (FTY720) administered orally once daily in patients with relapsing forms of multiple sclerosis ALTITUDE: A randomised, double-blind, placebo-controlled, parallel-group study to determine whether, in patients with type 2 diabeted at high risk for cardiovascular and renal events, aliskiren, on top of conventional treatment, reduces cardiovascular and renal events, aliskiren, on top of conventional treatment, reduces cardiovascular and renal morbidity and mortality FINISH 3: A Phase 3, Randomised, Single Blind, Controlled Trial of Topical Fibrocaps in Intraoperative Surgical Hemostasis GWCA0962: A double blind, randomized, placebo-controlled, parallel group study of Sativex oromucosal spray (Sativex®; Nabiximols) as adjunctive therapy in relieving uncontrolled persistent chronic pain in patients with	10/H1107/42 09/H0706/8 11/NE/0038 10/H0904/49 07/H0606/126	6 10 5 N a 8 S	31/08/2012 18/05/2012 02/05/2012 10 date greed with ponsor 01/01/2012	Up Complete Closed - Follow Up Complete	No No No
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration Of QUTENZA For The Treatment Of Peripheral Neuropathic Pain A Phase 3, Randomized, Double-Blind, Double-Dummy, Parallel Group, Multi-Center, Multi-National Study for Evaluation of Efficacy and Safety of DU-176b Versus Warfarin In Subjects with Atrial Fibrillation – Effective Anticoagulation with Factor Xa Next Generation in Atrial Fibrillation (ENGAGE – AF TIMI - 48) A Randomized, Double-Blind, Placebo-Controlled, Multicenter, Parallel-Group Study to Evaluate the Efficacy and Safety of Brivaracetam in Subjects (=16 To 80 Years Old) With Partial Onset Seizures A single arm, open-label, multicenter study evaluating the long-term safety and tolerability of 0.5 mg fingolimod (FTY720) administered orally once daily in patients with relapsing forms of multiple sclerosis ALTITUDE: A randomised, double-blind, placebo-controlled, parallel-group study to determine whether, in patients with type 2 diabeted at high risk for cardiovascular and renal events, aliskiren, on top of conventional treatment, reduces cardiovascular and renal morbidity and mortality FINISH 3: A Phase 3, Randomised, Single Blind, Controlled Trial of Topical Fibrocaps in Intraoperative Surgical Hemostasis GWCA0962: A double blind, randomized, placebo-controlled, parallel group study of Sativex oromucosal spray (Sativex®; Nabiximols) as adjunctive therapy in relieving uncontrolled persistent chronic pain in patients with advanced cancer, who experience inadequate analgesia during optimized	10/H1107/42 09/H0706/8 11/NE/0038 10/H0904/49 07/H0606/126 12/NE/0181	6 10 5 N a; 8 S	31/08/2012 18/05/2012 02/05/2012 10 date greed with ponsor 01/01/2012 31/03/2013	Up Complete Closed - Follow Up Complete	No No No No
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration Of QUTENZA For The Treatment Of Peripheral Neuropathic Pain A Phase 3, Randomized, Double-Blind, Double-Dummy, Parallel Group, Multi-Center, Multi-National Study for Evaluation of Efficacy and Safety of DU-176b Versus Warfarin In Subjects with Atrial Fibrillation – Effective Anticoagulation with Factor Xa Next Generation in Atrial Fibrillation (ENGAGE – AF TIMI - 48) A Randomized, Double-Blind, Placebo-Controlled, Multicenter, Parallel-Group Study to Evaluate the Efficacy and Safety of Brivaracetam in Subjects (=16 To 80 Years Old) With Partial Onset Seizures A single arm, open-label, multicenter study evaluating the long-term safety and tolerability of 0.5 mg fingolimod (FTY720) administered orally once daily in patients with relapsing forms of multiple sclerosis ALTITUDE: A randomised, double-blind, placebo-controlled, parallel-group study to determine whether, in patients with type 2 diabeted at high risk for cardiovascular and renal events, aliskiren, on top of conventional treatment, reduces cardiovascular and renal events, aliskiren, on top of conventional treatment, reduces cardiovascular and renal morbidity and mortality FINISH 3: A Phase 3, Randomised, Single Blind, Controlled Trial of Topical Fibrocaps in Intraoperative Surgical Hemostasis GWCA0962: A double blind, randomized, placebo-controlled, parallel group study of Sativex oromucosal spray (Sativex®; Nabiximols) as adjunctive therapy in relieving uncontrolled persistent chronic pain in patients with advanced cancer, who experience inadequate analgesia during optimized chronic opioid therapy	10/H1107/42 09/H0706/8 11/NE/0038 10/H0904/49 07/H0606/126	6 10 5 N a 8 S	31/08/2012 18/05/2012 02/05/2012 10 date greed with ponsor 01/01/2012 31/03/2013	Up Complete Closed - Follow Up Complete	No No No
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration Of QUTENZA For The Treatment Of Peripheral Neuropathic Pain A Phase 3, Randomized, Double-Blind, Double-Dummy, Parallel Group, Multi-Center, Multi-National Study for Evaluation of Efficacy and Safety of DU-176b Versus Warfarin In Subjects with Atrial Fibrillation – Effective Anticoagulation with Factor Xa Next Generation in Atrial Fibrillation (ENGAGE – AF TIMI - 48) A Randomized, Double-Blind, Placebo-Controlled, Multicenter, Parallel-Group Study to Evaluate the Efficacy and Safety of Brivaracetam in Subjects (=16 To 80 Years Old) With Partial Onset Seizures A single arm, open-label, multicenter study evaluating the long-term safety and tolerability of 0.5 mg fingolimod (FTY720) administered orally once daily in patients with relapsing forms of multiple sclerosis ALTITUDE: A randomised, double-blind, placebo-controlled, parallel-group study to determine whether, in patients with type 2 diabeted at high risk for cardiovascular and renal events, aliskiren, on top of conventional treatment, reduces cardiovascular and renal morbidity and mortality FINISH 3: A Phase 3, Randomised, Single Blind, Controlled Trial of Topical Fibrocaps in Intraoperative Surgical Hemostasis GWCA0962: A double blind, randomized, placebo-controlled, parallel group study of Sativex oromucosal spray (Sativex®; Nabiximols) as adjunctive therapy in relieving uncontrolled persistent chronic pain in patients with advanced cancer, who experience inadequate analgesia during optimized chronic opioid therapy A ClinIcal Evaluation of ST Changes in a Group of Patlents having Ventricular	10/H1107/42 09/H0706/8 11/NE/0038 10/H0904/49 07/H0606/126 12/NE/0181	6 10 5 N a; 8 S 10 10 10	31/08/2012 18/05/2012 02/05/2012 10 date greed with ponsor 01/01/2012 31/03/2013	Up Complete Closed - Follow Up Complete	No No No No No
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration Of QUTENZA For The Treatment Of Peripheral Neuropathic Pain A Phase 3, Randomized, Double-Blind, Double-Dummy, Parallel Group, Multi-Center, Multi-National Study for Evaluation of Efficacy and Safety of DU-176b Versus Warfarin In Subjects with Atrial Fibrillation – Effective Anticoagulation with Factor Xa Next Generation in Atrial Fibrillation (ENGAGE – AF TIMI - 48) A Randomized, Double-Blind, Placebo-Controlled, Multicenter, Parallel-Group Study to Evaluate the Efficacy and Safety of Brivaracetam in Subjects (=16 To 80 Years Old) With Partial Onset Seizures A single arm, open-label, multicenter study evaluating the long-term safety and tolerability of 0.5 mg fingolimod (FTY720) administered orally once daily in patients with relapsing forms of multiple sclerosis ALTITUDE: A randomised, double-blind, placebo-controlled, parallel-group study to determine whether, in patients with type 2 diabeted at high risk for cardiovascular and renal events, aliskiren, on top of conventional treatment, reduces cardiovascular and renal morbidity and mortality FINISH 3: A Phase 3, Randomised, Single Blind, Controlled Trial of Topical Fibrocaps in Intraoperative Surgical Hemostasis GWCA0962: A double blind, randomized, placebo-controlled, parallel group study of Sativex oromucosal spray (Sativex®; Nabiximols) as adjunctive therapy in relieving uncontrolled persistent chronic pain in patients with advanced cancer, who experience inadequate analgesia during optimized chronic opioid therapy A ClinIcal Evaluation of ST Changes in a Group of Patlents having Ventricular ArrHyThmias	10/H1107/42 09/H0706/8 11/NE/0038 10/H0904/49 07/H0606/126 12/NE/0181	6 10 5 N a; 8 S	31/08/2012 18/05/2012 02/05/2012 10 date greed with ponsor 01/01/2012 31/03/2013	Up Complete Closed - Follow Up Complete	No No No No
received prior anti-HER2 and chemotherapy-based treatment. A Multicentre, Single-Arm, Open-Label Study Of The Repeated Administration Of QUTENZA For The Treatment Of Peripheral Neuropathic Pain A Phase 3, Randomized, Double-Blind, Double-Dummy, Parallel Group, Multi-Center, Multi-National Study for Evaluation of Efficacy and Safety of DU-176b Versus Warfarin In Subjects with Atrial Fibrillation – Effective Anticoagulation with Factor Xa Next Generation in Atrial Fibrillation (ENGAGE – AF TIMI - 48) A Randomized, Double-Blind, Placebo-Controlled, Multicenter, Parallel-Group Study to Evaluate the Efficacy and Safety of Brivaracetam in Subjects (=16 To 80 Years Old) With Partial Onset Seizures A single arm, open-label, multicenter study evaluating the long-term safety and tolerability of 0.5 mg fingolimod (FTY720) administered orally once daily in patients with relapsing forms of multiple sclerosis ALTITUDE: A randomised, double-blind, placebo-controlled, parallel-group study to determine whether, in patients with type 2 diabeted at high risk for cardiovascular and renal events, aliskiren, on top of conventional treatment, reduces cardiovascular and renal morbidity and mortality FINISH 3: A Phase 3, Randomised, Single Blind, Controlled Trial of Topical Fibrocaps in Intraoperative Surgical Hemostasis GWCA0962: A double blind, randomized, placebo-controlled, parallel group study of Sativex oromucosal spray (Sativex®; Nabiximols) as adjunctive therapy in relieving uncontrolled persistent chronic pain in patients with advanced cancer, who experience inadequate analgesia during optimized chronic opioid therapy A Clinical Evaluation of ST Changes in a Group of Patlents having Ventricular	10/H1107/42 09/H0706/8 11/NE/0038 10/H0904/49 07/H0606/126 12/NE/0181	6 10 5 N a; 8 S 10 10 10	31/08/2012 18/05/2012 02/05/2012 10 date greed with ponsor 01/01/2012 31/03/2013	Up Complete Closed - Follow Up Complete In set up	No No No No No

Delivering Clinical Research June 2013

moderate hepatic impairment (Child-Pugh B) with genotype 1b chronic hepatitis C infection	13/LO/0268	23/11/2016 li	n set up	N/A
B1971009 Phase 3 study to assess Meningococcal B Vaccine	11/YH/0379	No date agreed with Sponsor I	n set up	N/A