


Physiotherapy exercises

following transtibial (below knee) amputation



Introduction

This sheet has been designed to help you remember the exercises that you have been taught by your physiotherapist. These exercises will help to keep you independent either in a wheelchair or using an artificial leg (prosthesis).

Exercise helps you to maintain your strength and mobility, and can also improve your blood flow.

You should try to continue with your exercises even after you have been discharged from treatment.

Remember, all exercises should be done slowly and smoothly. If you feel any pain, stop and report it to your physiotherapist or doctor.

Exercise 1: Static quadriceps

- Push your legs straight out in front of you
- Push the back of your knees into the bed and tighten the thigh muscles
- Hold for 5 seconds
- Repeat this ___ times



Exercise 2: Straight leg raise

- Extend your legs out in front of you
- Tighten the thigh muscles in your amputated leg
- Lift the leg off the bed
- Hold for ___ seconds
- Slowly lower
- Repeat ___ times

Repeat the above with the other leg.



Exercise 3: Inner range quads

- Sit with both legs out in front of you
- Place a rolled up towel, blanket or pillow under the knee of your amputated leg
- Straighten your knee
- Hold for 5 seconds
- Repeat ___ times

Repeat the above with the other leg.



Exercise 4: Hip adduction with resistance

- Sit with both legs out in front of you
- Place a pillow or rolled up towel between your knees
- Squeeze your legs together
- Hold for 5 seconds
- Repeat ___ times

NB This exercise can also be performed when sitting in a wheelchair or at the edge of a bed.



Exercise 5: Outer range quadriceps

- Sit on a chair or on the edge of the bed with your hands on your lap
- Straighten one knee
- Hold for 5 seconds
- Now bend your knee
- Repeat ___ times

Repeat the above with the other leg.



Exercise 6: Static gluteal contractions

- Lie on your back
- Keep both legs straight and close together
- Squeeze your buttocks as tightly as possible
- Hold for 5 seconds
- Repeat ___ times



Exercise 7: Hip flexor stretch

- Lie on your back, preferably without a pillow
- Bend your knee towards your chest and hold it there with your hands
- Push your opposite leg down flat on to the bed
- Hold for 30 – 60 seconds, then relax
- Repeat ___ times

Repeat the above with the other leg.



Exercise 8: Bridging

- Lie on your back with your arms at the side
- Place a couple of firm pillows or rolled up blankets under your thighs
- Pull in your stomach, tighten your buttocks and lift your bottom up off the bed
- Hold for 5 seconds
- Repeat ___ times


To make this exercise more difficult, place your arms across your chest as shown in the picture.



Exercise 9: Hip flexion and extension in side lying

- Lie on your side
- Bend the bottom leg
- Lift your top leg slightly
- Bend your knee fully towards your chest
- Straighten your knee and push your leg backwards
- Repeat ___ times

NB Try not to let your hips roll forwards or backwards. Repeat the above with the other leg.



Exercise 10: Hip abduction in side lying

- Lie on your side
- Bend the bottom leg
- Keep your hips and top leg in line with your body
- Slowly lift your top leg up, keeping your knee straight
- Slowly lower
- Repeat ___ times

NB Try not to let your hips roll forwards or backwards.

Repeat the above with the other leg.



Exercise 11: Knee flexion in prone lying

- Lie on your stomach
- Place your arms in a comfortable position
- Keeping your thigh on the bed, bend your knee as far as possible
- Hold for 5 seconds
- Straighten your knee
- Repeat ___ times

Repeat the above with the other leg.



Amputee therapy team
Rehabilitation Gymnasium
Douglas Bader Rehabilitation Centre
Queen Mary's Hospital
Roehampton
London
SW15 5PN

Tel: 020 8487 6042



Reference: THE_ETA_01

Published: July 2017

Review date: July 2019

Job Ref: 101680

Designed and produced by Media Services, SGUL