

Department of Renal Medicine and Transplantation


Removal of a Ureteric Stent following a Kidney Transplant

This information is for patients who have recently had a kidney transplant and who require a flexible cystoscopy and removal of the ureteric stent. This explains what happens before, during and after the procedure. It also tells you about the risks and side effects involved.

If you have any further questions, please speak to a doctor or nurse caring for you.

What is a ureteric stent?

The ureter is the tube that links the kidney to the bladder. A stent is a plastic tube that is inserted into the ureter at the time of your kidney transplant. It helps to drain urine from the new transplant into your bladder and allow healing. It helps to reduce the risk of urine leak at the new join between the ureter and the bladder. It also helps to prevent narrowing of the ureter where it joins the bladder, which may cause a blockage.



Why does it need to be removed?

- The stent is not needed after two to four weeks, as all the tissues should be healed. It can occasionally be left longer if advised by your transplant doctor or surgeon.

- Leaving the stent in for too long will increase the risk of urine infection or possible stone formation. These conditions may affect your transplant function.

Type of stents


We have two types of stents. These stents are similar except one has a magnet attached to one end of the stent. Depending on various factors, the implanting surgeon will decide which stent is best to place inside the transplant kidney ureter. The magnetic stent can be removed in outpatient clinic; however for a non-magnetic stent patients have to go to theatre and have flexible cystoscopy.

What is a magnetic stent?

The magnetic stent has a magnet attached to one end which goes into the urinary bladder. The benefit of it is that it can be taken out in outpatient clinic without a need to go to theatre.

How is a magnetic stent removed?

The magnetic stent comes with a removal probe with a magnet at one end of the probe. The probe is passed through the urethra into the urinary bladder. Both magnets click together and the stent is then pulled out.



What is a flexible cystoscopy?

This is an examination of the inside of your bladder using a fine, soft telescopic tube (the flexible cystoscope). The cystoscope is inserted through the urethra into the urinary bladder so that the doctor can see and remove the non-magnetic stent. In a majority of cases this procedure is done under local anaesthesia as a day case procedure.

What are the risks?

As with any surgical procedure there is a small chance of side-effects or complications such as:

- Infection
- Bleeding
- Risk of damaging the urethra or bladder that may need surgery. This risk is very low.
- Risk of partial or complete blockage of ureter that may need further management. This risk is also very low.

Contact us if you experience:

- Increasing pain or discomfort over your transplanted kidney or bladder area
- A high temperature
- Mild bleeding that does not resolve after two days or if you are passing blood clots at any time.

How can I prepare for removal of a ureteric stent?

- You can take your normal medications.
- You can eat and drink normally unless you have a peritoneal dialysis tube. This may be removed at the same time as the stent and you may require sedation or general anaesthesia.
- You will be given an antibiotic tablet before the procedure.

- The procedure will be explained to you and you will be asked to sign a consent form.

Asking for your consent

It is important that you feel involved in decisions about your care. For some treatments, you will be asked to sign a consent form to say that you agree to have the treatment and understand what it involves. You can withdraw your consent at any time, even if you have said 'yes' previously. If you would like more details about our consent process, please ask for a copy of our policy.

What happens during removal of a ureteric stent?

- The procedure will be performed in a clean outpatient room or operation theatre.
- The doctor will clean the surrounding area of skin with a mild disinfectant.
- Local anaesthetic gel is applied to numb the area, which can be mildly uncomfortable. This takes few minutes to work.
- For a magnetic stent a probe with a magnetic end is gently inserted into the urinary bladder through the urethra.
- For the non-magnetic stent the cystoscope is gently inserted into the urethra and up into the bladder.
- Men may be asked to cough or try to pass urine to allow the cystoscope to enter the bladder.
- As the stent is removed, you may feel like emptying your bladder.
- The procedure takes only few minutes.

Will I feel any pain?

You may feel some stinging as the magnetic probe or cystoscope enters the bladder.

What happens after the stent removal?

- You will be returned to the clinic or ward.
- You will be asked to empty your bladder.
- You may notice some blood in your urine but this should not concern you or delay you going home. Usually, it settles down very quickly.
- You may feel a little burning pain when you pass urine; this will get better after a day or so.
- After discharge from hospital, you can return to your normal activities.

Will I have a follow-up appointment?

You should have a clinic appointment within the next few days to monitor your kidney function and check for any issues.

Contact us

Transplant clinic on 020 8725 2455 (Between 8am and 4pm).
Champneys ward on 020 8725 0062 or 020 8725 1080 (Out of hours).

For more information leaflets on conditions, procedures, treatments and services offered at our hospitals, please visit www.stgeorges.nhs.uk

Additional services

Patient Advice and Liaison Service (PALS)

PALS can offer you advice and information when you have comments or concerns about our services or care. You can contact the PALS team on the advisory telephone line Monday, Tuesday, Thursday and Friday from 2pm to 5pm.

A Walk-in service is available:

Monday, Tuesday and Thursday between 10am and 4pm

Friday between 10am and 2pm.

Please contact PALS in advance to check if there are any changes to opening times.

The Walk-in and Advisory telephone services are closed on Wednesdays.

PALS is based within the hospital in the ground floor main corridor between Grosvenor and Lanesborough Wing.

Tel: 020 8725 2453 **Email:** pals@stgeorges.nhs.uk

NHS UK

The NHS provides online information and guidance on all aspects of health and healthcare, to help you make decisions about your health. **Web:** www.nhs.uk

NHS 111

You can call 111 when you need medical help fast but it's not a 999 emergency. NHS 111 is available 24 hours a day, 365 days a year. Calls are free from landlines and mobile phones. **Tel:** 111

