

Metabolic Dysfunction Associated Steatotic Liver Disease (MASLD)

This leaflet offers information about Metabolic Dysfunction Associated Steatotic Liver Disease (MASLD). If you have any further questions or concerns about MASLD, please speak to the staff member in charge of your care.


What is MASLD and why have I got it?

MASLD is the name given to a condition in which fat accumulates in the liver in people who drink little or no alcohol. MASLD is associated with being overweight or obese and / or having diabetes. It is also linked to high cholesterol levels and high blood pressure. Most people with MASLD do not develop serious liver problems but in some people, the build-up of fat in the liver can lead to serious liver disease (cirrhosis). People with MASLD also have an increased risk of having a heart attack and stroke. MASLD is a common problem, affecting more than a third of the population in the UK but the good news is that it is reversible once identified.

What are the signs and symptoms?

There are often no symptoms with MASLD although an enlarged fatty liver can cause mild pain in the right upper side of the abdomen. If the liver starts to fail yellowing of the eyes, confusion, abdominal swelling due to a build-up of fluid in the abdomen and spontaneous bleeding and bruising can occur, although these are rare symptoms. Most commonly you will not have any symptoms and MASLD is diagnosed when you have blood tests carried out that detect abnormal liver tests or when fat in the liver is seen on an ultrasound scan of the liver. This ultrasound scan may be undertaken because you have risk factors for MASLD or the condition may be found when you have a scan for another reason.

Stages of Liver Disease



Do I need any tests to confirm the diagnosis?

When abnormal liver tests are detected on your blood tests it is important to ensure that there is no other cause for the abnormal liver tests so you will need to have further blood tests carried out to look for other causes, e.g. viral and immune causes. An ultrasound scan of the liver will also be carried out to look for fat within the liver and to ensure there is no other cause for the abnormal liver tests. A 'fibrosis score' can then be calculated (a score using blood tests and your age) that looks at your risk of developing scarring in the liver and whether you can be looked after by your GP or need to be seen in the Liver Clinic at St George's Hospital.

Is there any treatment for MASLD?

There are currently no effective drug treatments for MASLD although there is a lot of research being undertaken in this area. If you would be interested in taking part in a research project in MASLD at St George's Hospital please do let our Hepatology Clinical Assessment Nurse know.

Is there anything I can do to help myself?

Currently the best way to treat MASLD is through reversing the underlying cause. This includes weight loss if you are overweight (a sustained 10% reduction in body weight can reverse the fatty liver and prevent damage developing), with both a low fat diet and a mediterranean diet helping reduce fat in the liver. Regular moderate exercise (30 minutes 3-5 times a week), so you feel the effort and get a little warm and sweaty but not too out of breath, helps burn fat in the liver. If you have diabetes it is vital you get good control of your blood sugar because high blood sugar levels can cause more fat to be laid down in the liver. Avoid drinking alcohol as it causes a fatty liver and is high in calories.



Useful sources of information

British Liver Trust

[MASLD, NAFLD and fatty liver disease - British Liver Trust](#)

NHS

<https://www.nhs.uk/conditions/non-alcoholic-fatty-liver-disease>

<https://www.nhs.uk/live-well>

Contact us

If you have any questions or concerns about MASLD, please contact the hepatology team on 020 8725 7158 (Monday to Friday, 10am to 4pm).

For more information leaflets on conditions, procedures, treatments and services offered at our hospitals, please visit www.stgeorges.nhs.uk

Additional services

Patient Advice and Liaison Service (PALS)

PALS can offer you advice and information when you have comments or concerns about our services or care. You can contact the PALS team on the advisory telephone line Monday, Tuesday, Thursday and Friday from 2pm to 5pm.

A Walk-in service is available:

Monday, Tuesday and Thursday between 10am and 4pm.

Friday between 10am and 2pm.

Please contact PALS in advance to check if there are any changes to opening times.

The Walk-in and Advisory telephone services are closed on Wednesdays.

PALS is based within the hospital in the ground floor main corridor between Grosvenor and Lanesborough Wing.

Tel: 020 8725 2453 **Email:** pals@stgeorges.nhs.uk

NHS UK

The NHS provides online information and guidance on all aspects of health and healthcare, to help you make decisions about your health. **Web:** www.nhs.uk

NHS 111

You can call 111 when you need medical help fast but it's not a 999 emergency. NHS 111 is available 24 hours a day, 365 days a year. Calls are free from landlines and mobile phones. **Tel:** 111

AccessAble

You can download accessibility guides for all our services by searching 'St George's Hospital' on the AccessAble website (www.accessable.co.uk). The guides are designed to ensure everyone – including those with accessibility needs – can access our hospital and community sites with confidence.



Reference: GST_MASLD_02 **Published:** August 2024 **Review date:** August 2026